

# Citrex

Diverse in Nature



The Organic Response to:

- ▶ **Poultry Pathogens**
- ▶ **Respiratory Disease**
- ▶ **Salmonellosis**



**LIQUID**

## USES & DOSAGES



### POULTRY DRINKING WATER

Substitute of promoters 0.05 to 0.1 ml / lt continuously  
 Stress situation 0.2 ml / lt during and 3 days after stress  
 Salmonella positive feces 0.1 ml / lt continuously for 21 days  
 Coadjutant to diarrhea 0.2 to 0.3 ml / lt

### COADJUTANT AEROSOL / SPRAY TREATMENT

Respiratory problems viral or bacterial. Dosage per 250 m<sup>2</sup>:

**Preventive** 1.0 ml / lt 3 times per week  
**Coadjutant to treatments:** 1.0 ml / lt 2 per day for 4 to 6 days

### DISINFECTION

General: 0.1 to 0.2 ml/lt Hatcheries: 0.4 to 0.6 ml/lt Fertile eggs: 0.8 to 1.0 ml / lt

## Citrex AGAINST MICROORGANISMS

### BACTERIOSTATIC (MIC)

BACTERIA	ppm
<i>Bordetella avium</i>	50
<i>Escherichia coli K-88</i>	100
<i>Enterococcus aerogenes</i>	50
<i>Enterococcus faecalis</i>	25
<i>Listeria monocytogenes</i>	12.5
<i>Pasteurella multocida</i>	25
<i>Proteus vulgaris</i>	25
<i>Pseudomona aeruginosa</i>	200
<i>Salmonella choleraesuis</i>	50
<i>Salmonella typhimurium</i>	50
<i>Staphylococcus aureus</i>	12.5
<i>Streptococcus suis sero 2</i>	13
<i>Vibrio vulnificus</i>	75
<i>Vibrio parahaemolyticus</i>	50

### FUNGISTATIC (MIC)

FUNGI	ppm
<i>Aspergillus niger</i>	50
<i>Candida albicans</i>	6.25
<i>Trichophyton mentagrophyte</i>	25

### BACTERICIDAL

BACTERIA	ppm
<i>Escherichia coli C-68</i>	100
<i>Escherichia coli C-73</i>	200
<i>Salmonella enteritidis</i>	200
<i>Salmonella typhimurium</i>	200
<i>Salmonella seftemberg</i>	200
<i>Pseudomona C-102</i>	600
<i>Pseudomona C-107</i>	400

### VIRICIDAL

VIRUS	ppm
<i>Laringotraqueitis</i>	100
<i>Avian Influenza</i>	100
<i>Infectious Bursal Disease</i>	100
<i>Chicken Pox</i>	100
<i>Infectious Bronchitis</i>	100
<i>Newcastle Disease</i>	100
<i>Pseudorabies</i>	200
<i>PRRS</i>	200
<i>Swine Fever</i>	400